

ZNF391 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13177**Specification**

ZNF391 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9UJN7
Other Accession	NM_001076781 , NP_001070249
Reactivity	Human, Bovine, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

ZNF391 antibody - N-terminal region - Additional Information**Gene ID** 346157

Alias Symbol	dj153G14.3
Other Names	
Zinc finger protein 391, ZNF391	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF391 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF391 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF391 antibody - N-terminal region - Protein Information**Name** ZNF391**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF391 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF391 antibody - N-terminal region - Images



WB Suggested Anti-ZNF391 Antibody Titration: 0.2-1 µg/ml

Positive Control: 721_B cell lysate

ZNF391 is supported by BioGPS gene expression data to be expressed in 721_B

ZNF391 antibody - N-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).

Mungall A.J., et al. Nature 425:805-811(2003).

Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.