

SCRT2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13214**Specification**

SCRT2 antibody - middle region - Product Information

Application	WB
Primary Accession	O9NQ03
Other Accession	NM_033129 , NP_149120
Reactivity	Human, Rat, Yeast, Guinea Pig, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

SCRT2 antibody - middle region - Additional Information**Gene ID** 85508**Alias Symbol** **ZNF898B****Other Names**

Transcriptional repressor scratch 2, Scratch homolog 2 zinc finger protein, SCRT2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SCRT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SCRT2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

SCRT2 antibody - middle region - Protein Information**Name** SCRT2**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

SCRT2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCRT2 antibody - middle region - Images



WB Suggested Anti-SCRT2 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Human Muscle

SCRT2 antibody - middle region - References

Wan D.F., et al. Submitted (JUL-2002) to the EMBL/GenBank/DDBJ databases.

Deloukas P., et al. Nature 414:865-871(2001).