

ZNF280A antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13216**Specification**

ZNF280A antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P59817
Other Accession	NM_080740 , NP_542778
Reactivity	Human, Rat, Guinea Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60kDa KDa

ZNF280A antibody - C-terminal region - Additional Information**Gene ID** 129025**Alias Symbol** 3'OY11.1, SUHW1, ZNF280, ZNF636**Other Names**

Zinc finger protein 280A, 3'OY11.1, Suppressor of hairy wing homolog 1, Zinc finger protein 636, ZNF280A, SUHW1, ZNF280, ZNF636

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF280A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF280A antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF280A antibody - C-terminal region - Protein Information**Name** ZNF280A**Synonyms** SUHW1, ZNF280, ZNF636**Function**

May function as a transcription factor.

Cellular Location


Nucleus.

ZNF280A antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF280A antibody - C-terminal region - Images



WB Suggested Anti-ZNF280A Antibody Titration: 0.2-1 μ g/ml

Positive Control: Human brain

ZNF280A antibody - C-terminal region - References

Kawasaki K., et al. Genome Res. 7:250-261(1997).

Dunham I., et al. Nature 402:489-495(1999).