

ZNF790 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13248**Specification**

ZNF790 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6PG37
Other Accession	NM_206894 , NP_996777
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74kDa KDa

ZNF790 antibody - N-terminal region - Additional Information**Gene ID** 388536

Alias Symbol	FLJ20350, MGC62100
Other Names	
Zinc finger protein 790, ZNF790	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF790 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF790 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF790 antibody - N-terminal region - Protein Information**Name** ZNF790**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

ZNF790 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF790 antibody - N-terminal region - Images



WB Suggested Anti-ZNF790 Antibody Titration: 0.2-1 μ g/ml
Positive Control: 721_B cell lysate

ZNF790 antibody - N-terminal region - References

Grimwood J., et al. Nature 428:529-535(2004).