

ZNF320 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13249**Specification**

ZNF320 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A2RRD8
Other Accession	NM_207333 , NP_997216
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59kDa KDa

ZNF320 antibody - N-terminal region - Additional Information**Gene ID** 162967

Alias Symbol	DKFZp686G16228, ZFPL
Other Names	
Zinc finger protein 320, ZNF320	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF320 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF320 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF320 antibody - N-terminal region - Protein Information**Name** ZNF320**Function**

May be involved in transcriptional regulation.

Cellular Location

Nucleus.

Tissue Location


Expressed at high levels in placenta, pancreas, kidney and heart, and at lower levels in liver, skeletal muscle and brain.

ZNF320 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF320 antibody - N-terminal region - Images



WB Suggested Anti-ZNF320 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: MCF7 cell lysate

ZNF320 antibody - N-terminal region - References

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Chernova O.B.,et al.Oncogene 20:5378-5392(2001).