

hCG_1660138 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13258**Specification**

hCG_1660138 antibody - N-terminal region - Product Information

Application	WB
Other Accession	XM_066752 , XP_066752
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31kDa KDa

hCG_1660138 antibody - N-terminal region - Additional Information

Alias Symbol **LOC139542, hCG_1660138**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-hCG_1660138 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


hCG_1660138 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

hCG_1660138 antibody - N-terminal region - Protein Information**hCG_1660138 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

hCG_1660138 antibody - N-terminal region - Images



WB Suggested Anti-hCG_1660138 Antibody Titration: 0.2-1 μ g/ml
Positive Control: 721_B cell lysate