

**PEG10 Antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13294****Specification**

---

**PEG10 Antibody - C-terminal region - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB  |
| Primary Accession | <a href="#">Q86TG7</a>                                      |
| Other Accession   | <a href="#">NM_001172438</a> , <a href="#">NP_001165909</a> |
| Reactivity        | Human, Mouse, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog   |
| Predicted         | Human, Mouse, Rabbit, Pig, Horse, Bovine, Guinea Pig        |
| Host              | Rabbit  |
| Clonality         | Polyclonal  |
| Calculated MW     | 44kDa kDa   |

**PEG10 Antibody - C-terminal region - Additional Information****Gene ID** 23089**Alias Symbol** EDR, HB-1, MEF3L, Mar2, Mart2, RGAG3**Other Names**

Retrotransposon-derived protein PEG10, Embryonal carcinoma differentiation-regulated protein, Mammalian retrotransposon-derived protein 2, Myelin expression factor 3-like protein 1, MEF3-like protein 1, Paternally expressed gene 10 protein, Retrotransposon gag domain-containing protein 3, Retrotransposon-derived gag-like polyprotein, Ty3/Gypsy-like protein, PEG10, EDR, KIAA1051, MAR2, MART2, MEF3L1, RGAG3

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-PEG10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

PEG10 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**PEG10 Antibody - C-terminal region - Protein Information****Name** PEG10 {ECO:0000303|PubMed:11318613, ECO:0000312|HGNC:HGNC:14005}**Function**

Retrotransposon-derived protein that binds its own mRNA and self-assembles into virion-like capsids (PubMed:<a href="http://www.uniprot.org/citations/34413232" target="\_blank">34413232</a>). Forms virion-like extracellular vesicles that encapsulate their

own mRNA and are released from cells, enabling intercellular transfer of PEG10 mRNA (PubMed:<a href="http://www.uniprot.org/citations/34413232" target="\_blank">34413232</a>). Binds its own mRNA in the 5'-UTR region, in the region near the boundary between the nucleocapsid (NC) and protease (PRO) coding sequences and in the beginning of the 3'-UTR region (PubMed:<a href="http://www.uniprot.org/citations/34413232" target="\_blank">34413232</a>). Involved in placenta formation: required for trophoblast stem cells differentiation (By similarity). Involved at the immediate early stage of adipocyte differentiation (By similarity). Overexpressed in many cancers and enhances tumor progression: promotes cell proliferation by driving cell cycle progression from G0/G1 (PubMed:<a href="http://www.uniprot.org/citations/12810624" target="\_blank">12810624</a>, PubMed:<a href="http://www.uniprot.org/citations/16423995" target="\_blank">16423995</a>, PubMed:<a href="http://www.uniprot.org/citations/26235627" target="\_blank">26235627</a>, PubMed:<a href="http://www.uniprot.org/citations/28193232" target="\_blank">28193232</a>). Enhances cancer progression by inhibiting the TGF-beta signaling, possibly via interaction with the TGF-beta receptor ACVRL1 (PubMed:<a href="http://www.uniprot.org/citations/15611116" target="\_blank">15611116</a>, PubMed:<a href="http://www.uniprot.org/citations/26235627" target="\_blank">26235627</a>, PubMed:<a href="http://www.uniprot.org/citations/30094509" target="\_blank">30094509</a>). May bind to the 5'-GCCTGTCTTT-3' DNA sequence of the MB1 domain in the myelin basic protein (MBP) promoter; additional evidences are however required to confirm this result (By similarity).

### Cellular Location

Extracellular vesicle membrane. Cytoplasm. Nucleus Note=Forms virion-like extracellular vesicles that are released from cells (PubMed:34413232). Detected predominantly in the cytoplasm of breast and prostate carcinomas, in hepatocellular carcinoma (HCC) and B-cell chronic lymphocytic leukemia (B-CLL) cells and in the Hep-G2 cell line (PubMed:12810624).

### Tissue Location

Expressed in the cytotrophoblast layer but not in the overlying syncytiotrophoblast of the placenta. Expressed in prostate and breast carcinomas but not in normal breast and prostate epithelial cells. Expressed in the Hep-G2 cell line (at protein level) Expressed in brain, liver, spleen, kidney, thymus, lung, ovary, testis, reactive lymph node, skeletal muscle, adipose tissue and placenta Expressed in pancreatic and hepatocellular carcinomas (HCC)


## PEG10 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PEG10 Antibody - C-terminal region - Images





Host: Rabbit  
Target Name: PEG10  
Sample Tissue: Thyroid Tumor lysates  
Antibody Dilution: 1.0 $\mu$ g/ml