

CCDC19 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13298**Specification**

CCDC19 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5VU18
Other Accession	NM_012337 , NP_036469
Reactivity	Human, Rat, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted Host	Pig, Horse, Bovine, Guinea Pig, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 66kDa KDa

CCDC19 antibody - N-terminal region - Additional Information

Alias Symbol **NESG1, RP11-190A12.6**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CCDC19 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CCDC19 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CCDC19 antibody - N-terminal region - Protein Information**CCDC19 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC19 antibody - N-terminal region - Images



WB Suggested Anti-CCDC19 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: 293T cell lysate