

SSX8 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13317**Specification**

SSX8 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q7RTT4
Other Accession	NM_174961 , NP_777621
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

SSX8 antibody - C-terminal region - Additional Information**Other Names**

Protein SSX8, SSX8

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SSX8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SSX8 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SSX8 antibody - C-terminal region - Protein Information**Name** SSX8P ([HGNC:19654](#))**Synonyms** SSX8**Function**

Could act as a modulator of transcription.

Tissue Location


Not detected in any normal or tumor tissues.

SSX8 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SSX8 antibody - C-terminal region - Images



WB Suggested Anti-SSX8 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Human Liver

SSX8 antibody - C-terminal region - References

Ross M.T., et al. Nature 434:325-337(2005).
Gure A.O., et al. Int. J. Cancer 101:448-453(2002).