

C8orf77 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13319**Specification**

C8orf77 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q0IIN9
Other Accession	NM_178535 , NP_848630
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23kDa KDa

C8orf77 antibody - C-terminal region - Additional Information

Alias Symbol **MGC141859, C8orf77**

Other Names

Putative uncharacterized protein ZNF252P-AS1, ZNF252P antisense RNA 1, ZNF252P antisense gene protein 1, ZNF252P-AS1, C8orf77

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C8orf77 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C8orf77 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

C8orf77 antibody - C-terminal region - Protein Information

Name ZNF252P-AS1

Synonyms C8orf77


C8orf77 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C8orf77 antibody - C-terminal region - Images



WB Suggested Anti-C8orf77 Antibody Titration: 0.2-1 µg/ml
Positive Control: 721_B cell lysate

C8orf77 antibody - C-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Nusbaum C., et al. Nature 439:331-335(2006).