

CROCCL2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13323**Specification**

CROCCL2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8IVE0
Other Accession	XM_057040 , XP_057040
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Bovine, Guinea Pig, Dog
Predicted	Human, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

CROCCL2 antibody - N-terminal region - Additional Information

Alias Symbol	KIAA0445, KIAA1922, dj798A10.2, CROCCL2
--------------	---

Other Names

Putative ciliary rootlet coiled-coil protein-like 2 protein, Ciliary rootlet coiled-coil protein pseudogene 3, CROCCP3, CROCCL2, KIAA1922

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CROCCL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CROCCL2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CROCCL2 antibody - N-terminal region - Protein Information

Name CROCCP3

Synonyms CROCCL2, KIAA1922


CROCCL2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CROCCL2 antibody - N-terminal region - Images



WB Suggested Anti-CROCCL2 Antibody Titration: 0.2-1 µg/ml
Positive Control: MCF7 cell lysate

CROCCL2 antibody - N-terminal region - References

Nagase T., et al. DNA Res. 8:179-187(2001).
Gregory S.G., et al. Nature 441:315-321(2006).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.