

NET1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13372**Specification**

NET1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5SQI7
Other Accession	NM_005863 , NP_005854
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62kDa KDa

NET1 antibody - N-terminal region - Additional Information

Alias Symbol **ARHGEF8, NET1A**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NET1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NET1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NET1 antibody - N-terminal region - Protein Information**NET1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NET1 antibody - N-terminal region - Images



WB Suggested Anti-NET1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: 293T cell lysate

NET1 is strongly supported by BioGPS gene expression data to be expressed in Human HEK293T cells

NET1 antibody - N-terminal region - Citations

- [The guanine nucleotide exchange factor Net1 facilitates the specification of dorsal cell fates in zebrafish embryos by promoting maternal \$\beta\$ -catenin activation.](#)