

RFPL2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13385**Specification**

RFPL2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O75678
Other Accession	NM_001098527 , NP_001091997
Reactivity	Human, Rabbit
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

RFPL2 antibody - C-terminal region - Additional Information**Gene ID** 10739

Alias Symbol	RNF79
Other Names	
Ret finger protein-like 2, RING finger protein 79, RFPL2, RNF79	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RFPL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RFPL2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RFPL2 antibody - C-terminal region - Protein Information**Name** RFPL2**Synonyms** RNF79**Tissue Location**


Seems to be expressed in prostate and less abundantly in adult brain, fetal liver, and fetal kidney

RFPL2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RFPL2 antibody - C-terminal region - Images



WB Suggested Anti-RFPL2 Antibody Titration: 0.2-1 μ g/ml

Positive Control: Human Liver

RFPL2 antibody - C-terminal region - References

- Seroussi E., et al. Genome Res. 9:803-814(1999).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004).
Dunham I., et al. Nature 402:489-495(1999).