

**Katna1 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13395****Specification**

---

**Katna1 antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O9WV86</a>
Other Accession	<a href="#">NM_011835</a> , <a href="#">NP_035965</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

**Katna1 antibody - C-terminal region - Additional Information****Other Names**

Katanin p60 ATPase-containing subunit A1 {ECO:0000255|HAMAP-Rule:MF\_03023}, Katanin p60 subunit A1 {ECO:0000255|HAMAP-Rule:MF\_03023}, 3.6.4.3 {ECO:0000255|HAMAP-Rule:MF\_03023}, Lipotransin, p60 katanin {ECO:0000255|HAMAP-Rule:MF\_03023}, Katna1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Katna1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Katna1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Katna1 antibody - C-terminal region - Protein Information**

**Name** Katna1

**Function**

Catalytic subunit of a complex which severs microtubules in an ATP-dependent manner. Microtubule severing may promote rapid reorganization of cellular microtubule arrays and the release of microtubules from the centrosome following nucleation. Microtubule release from the mitotic spindle poles may allow depolymerization of the microtubule end proximal to the spindle pole, leading to poleward microtubule flux and poleward motion of chromosome. The function in regulating microtubule dynamics at spindle poles seems to depend on the association of the katanin KATNA1:KATNB1 complex with ASPM which recruits it to microtubules. Reversely

KATNA1:KATNB1 can enhance ASPM blocking activity on microtubule minus-end growth. Microtubule release within the cell body of neurons may be required for their transport into neuronal processes by microtubule-dependent motor proteins. This transport is required for axonal growth.

### Cellular Location


Cytoplasm {ECO:0000255|HAMAP-Rule:MF\_03023}. Midbody {ECO:0000255|HAMAP-Rule:MF\_03023}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000255|HAMAP-Rule:MF\_03023}. Cytoplasm, cytoskeleton, spindle pole {ECO:0000255|HAMAP-Rule:MF\_03023}. Cytoplasm, cytoskeleton, spindle {ECO:0000250|UniProtKB:O75449}. Note=Predominantly cytoplasmic Localized diffusely in the cytoplasm during the interphase. During metaphase is localized throughout the cell and more widely dispersed than the microtubules. In anaphase and telophase is localized at the midbody region. Also localized to the interphase centrosome and the mitotic spindle poles. Enhanced recruitment to the mitotic spindle poles requires microtubules and interaction with KATNB1 (By similarity). Localizes within the cytoplasm, partially overlapping with microtubules, in interphase and to the mitotic spindle and spindle poles during mitosis (By similarity). {ECO:0000250|UniProtKB:O75449, ECO:0000255|HAMAP-Rule:MF\_03023}

### Katna1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Katna1 antibody - C-terminal region - Images



WB Suggested Anti-Katna1 Antibody Titration: 1.0 µg/ml  
Positive Control: Mouse Spleen

### Katna1 antibody - C-terminal region - References

Syu L.-J., et al. Mol. Cell 4:109-115(1999).  
Karabay A., et al. J. Neurosci. 24:5778-5788(2004).