

Crygs antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13406**Specification**

Crygs antibody - C-terminal region - Product Information

Application	WB
Primary Accession	POC5E9
Other Accession	NM_001109553 , NP_001103023
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19kDa KDa

Crygs antibody - C-terminal region - Additional Information**Gene ID** 689897**Other Names**

Beta-crystallin S, Gamma-S-crystallin, Gamma-crystallin S, Crygs

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Crygs antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Crygs antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Crygs antibody - C-terminal region - Protein Information**Name** Crygs**Function**


Crystallins are the dominant structural components of the vertebrate eye lens.

Crygs antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Crygs antibody - C-terminal region - Images



Host: Rabbit
Target Name: Crygs
Antibody Dilution: 1.0µg/ml
Sample Tissue: Rat Testis

Crygs antibody - C-terminal region - References

Gibbs R.A., et al. Nature 428:493-521(2004).