

Olfm3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13421**Specification**

Olfm3 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P63056
Other Accession	NM_153157 , NP_694797
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

Olfm3 antibody - C-terminal region - Additional Information**Gene ID** 229759

Alias Symbol	B230206G02Rik
Other Names	
Noelin-3, Olfactomedin-3, Optimedin, Olfm3, Noe3	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Olfm3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Olfm3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Olfm3 antibody - C-terminal region - Protein Information**Name** Olfm3**Synonyms** Noe3**Cellular Location**

Secreted. Synapse. Note=Isoform 2 is secreted more efficiently than isoform 1

Tissue Location


Expressed in the brain (at protein level).

Olfm3 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Olfm3 antibody - C-terminal region - Images



WB Suggested Anti-Olfm3 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Liver

Olfm3 antibody - C-terminal region - References

Torrado M., et al. Hum. Mol. Genet. 11:1291-1301(2002).
Carninci P., et al. Science 309:1559-1563(2005).
Schwenk J., et al. Neuron 74:621-633(2012).