

ACTRT2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13422**Specification**

ACTRT2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8TDY3
Other Accession	NM_080431 , NP_536356
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Mouse, Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

ACTRT2 antibody - C-terminal region - Additional Information**Gene ID** 140625**Alias Symbol** **ARPM2, ARPT2, Arp-T2, FLJ25424, HARPM2**
Other Names
Actin-related protein T2, ARP-T2, Actin-related protein M2, ACTRT2, ARPM2**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ACTRT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ACTRT2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ACTRT2 antibody - C-terminal region - Protein Information**Name** ACTRT2**Synonyms** ARPM2**Cellular Location**


Cytoplasm, cytoskeleton.

ACTRT2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACTRT2 antibody - C-terminal region - Images



WB Suggested Anti-ACTRT2 Antibody Titration: 0.2-1 µg/ml

Positive Control: Human Muscle

ACTRT2 antibody - C-terminal region - References

- Harata M., et al. Biochim. Biophys. Acta 1522:130-133(2001).
Heid H.W., et al. Exp. Cell Res. 279:177-187(2002).
Gregory S.G., et al. Nature 441:315-321(2006).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.