

SPATA17 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13443**Specification**

SPATA17 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O96L03
Other Accession	NM_138796 , NP_620151
Reactivity	Human, Rabbit
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

SPATA17 antibody - C-terminal region - Additional Information**Gene ID** 128153

Alias Symbol	IQCH, MSRG-11, MSRG11, RP11-144C20.1
Other Names	
Spermatogenesis-associated protein 17, SPATA17	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SPATA17 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SPATA17 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


SPATA17 antibody - C-terminal region - Protein Information**Name** SPATA17**Cellular Location**
Cytoplasm.**SPATA17 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPATA17 antibody - C-terminal region - Images



WB Suggested Anti-SPATA17 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Human brain

SPATA17 antibody - C-terminal region - References

Deng Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.