

TIGD4 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13468**Specification**

TIGD4 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8IY51
Other Accession	NM_145720 , NP_663772
Reactivity	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

TIGD4 antibody - N-terminal region - Additional Information**Gene ID** 201798**Alias Symbol** MGC43837**Other Names**

Tigger transposable element-derived protein 4, TIGD4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TIGD4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TIGD4 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TIGD4 antibody - N-terminal region - Protein Information**Name** TIGD4**Cellular Location**


Nucleus.

TIGD4 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TIGD4 antibody - N-terminal region - Images



WB Suggested Anti-TIGD4 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

TIGD4 antibody - N-terminal region - References

Hillier L.W., et al. Nature 434:724-731(2005).

Ota T., et al. Nat. Genet. 36:40-45(2004).