

Ociad2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13486**Specification**

Ociad2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q9D8W7
Other Accession	NM_026950 , NP_081226
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

Ociad2 antibody - N-terminal region - Additional Information**Gene ID** 433904

Alias Symbol	1810027I20Rik, MGC51598
Other Names	
OCIA domain-containing protein 2, Ociad2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ociad2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ociad2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ociad2 antibody - N-terminal region - Protein Information**Name** Ociad2**Cellular Location**


Endosome.

Ociad2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ociad2 antibody - N-terminal region - Images



WB Suggested Anti-Ociad2 Antibody Titration: 1.0 µg/ml

Positive Control: Mouse Heart

Ociad2 antibody - N-terminal region - References

Carninci P., et al. Science 309:1559-1563(2005).

Rardin M.J., et al. Proc. Natl. Acad. Sci. U.S.A. 110:6601-6606(2013).