

FAM71A antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13521**Specification**

FAM71A antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8IYT1
Other Accession	NM_153606 , NP_705834
Reactivity	Human, Rabbit, Pig, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rabbit, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 63kDa KDa

FAM71A antibody - C-terminal region - Additional Information**Gene ID** 149647

Alias Symbol	FLJ32796, RP11-338C15.4
Other Names	
Protein FAM71A, FAM71A	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-FAM71A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

FAM71A antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FAM71A antibody - C-terminal region - Protein Information**Name** GARIN4 ([HGNC:26541](#))**Synonyms** FAM71A**Function**

RAB2B effector protein required for the compacted Golgi morphology, probably through interaction with small GTPase RAB2B.

Cellular Location


Golgi apparatus.

FAM71A antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM71A antibody - C-terminal region - Images



WB Suggested Anti-FAM71A Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

FAM71A antibody - C-terminal region - References

Gregory S.G., et al. Nature 441:315-321(2006).