

PDE12 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13542**Specification**

PDE12 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	F6T1Q0
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

PDE12 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PDE12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

PDE12 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PDE12 Antibody - C-terminal region - Protein Information**PDE12 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PDE12 Antibody - C-terminal region - Images



90 kDa__
65 kDa__
40 kDa__
29 kDa__
22 kDa__

Host: Rabbit
Target Name: PDE12
Sample Tissue: Ovary Tumor lysates
Antibody Dilution: 1.0 μ g/ml