

CMTM8 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13544**Specification**

CMTM8 antibody - middle region - Product Information

Application	WB
Primary Accession	Q8IZV2
Other Accession	NM_178868 , NP_849199
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rat, Horse, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 19kDa KDa

CMTM8 antibody - middle region - Additional Information**Gene ID** 152189**Alias Symbol** CKLFSF8, CKLFSF8-V2**Other Names**

CKLF-like MARVEL transmembrane domain-containing protein 8, Chemokine-like factor superfamily member 8, CMTM8, CKLFSF8

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CMTM8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CMTM8 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

CMTM8 antibody - middle region - Protein Information**Name** CMTM8**Synonyms** CKLFSF8**Cellular Location**

[Isoform 1]: Membrane; Multi-pass membrane protein

Tissue Location


Highly expressed in liver and pancreas.

CMTM8 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CMTM8 antibody - middle region - Images



WB Suggested Anti-CMTM8 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate

CMTM8 antibody - middle region - References

Han W., et al. Genomics 81:609-617(2003).
Li D., et al. Int. J. Biochem. Cell Biol. 39:2107-2119(2007).
Muzny D.M., et al. Nature 440:1194-1198(2006).