

**PHACTR3 antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13558****Specification**

---

**PHACTR3 antibody - C-terminal region - Product Information**

|                   |                                                                         |
|-------------------|-------------------------------------------------------------------------|
| Application       | WB                                                                      |
| Primary Accession | <a href="#">B1AN69</a>                                                  |
| Other Accession   | <a href="#">NM_183246</a> , <a href="#">NP_899069</a>                   |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog          |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                                  |
| Clonality         | Polyclonal                                                              |
| Calculated MW     | 51kDa KDa                                                               |

**PHACTR3 antibody - C-terminal region - Additional Information**

|              |                                                 |
|--------------|-------------------------------------------------|
| Alias Symbol | C20orf101, H17739, MGC117178, SCAPIN1, SCAPININ |
|--------------|-------------------------------------------------|

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-PHACTR3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


**Precautions**

PHACTR3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**PHACTR3 antibody - C-terminal region - Protein Information****PHACTR3 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PHACTR3 antibody - C-terminal region - Images**

WB Suggested Anti-PHACTR3 Antibody Titration: 0.2-1  $\mu$ g/ml  
Positive Control: HepG2 cell lysate