

SCCPDH antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13603**Specification**

SCCPDH antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NBX0
Other Accession	NM_016002 , NP_057086
Reactivity	Human, Rat, Horse, Guinea Pig, Dog
Predicted	Human, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

SCCPDH antibody - C-terminal region - Additional Information**Gene ID** 51097**Alias Symbol** CGI-49, FLJ43187, RP11-439E19.2, NET11**Other Names**

Saccharopine dehydrogenase-like oxidoreductase, 1.-.-., SCCPDH

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SCCPDH antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SCCPDH antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


SCCPDH antibody - C-terminal region - Protein Information**Name** SCCPDH**SCCPDH antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCCPDH antibody - C-terminal region - Images



WB Suggested Anti-SCCPDH Antibody Titration: 0.2-1 µg/ml
Positive Control: Human Muscle

SCCPDH antibody - C-terminal region - References

Lai C.-H., et al. *Genome Res.* 10:703-713(2000).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Gregory S.G., et al. *Nature* 441:315-321(2006).
Bienvenut W.V., et al. Submitted (JAN-2010) to UniProtKB.
Burkard T.R., et al. *BMC Syst. Biol.* 5:17-17(2011).