

C2orf42 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13608**Specification**

C2orf42 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O9NWW7
Other Accession	NM_017880 , NP_060350
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Pig, Chicken, Horse
Clonality	Rabbit
Calculated MW	Polyclonal 63kDa KDa

C2orf42 antibody - C-terminal region - Additional Information**Gene ID** 54980**Alias Symbol** FLJ20558**Other Names**

Uncharacterized protein C2orf42, C2orf42

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C2orf42 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C2orf42 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


C2orf42 antibody - C-terminal region - Protein Information**Name** C2orf42**C2orf42 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C2orf42 antibody - C-terminal region - Images



WB Suggested Anti-C2orf42 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate

C2orf42 antibody - C-terminal region - References

Wiemann S., et al. *Genome Res.* 11:422-435(2001).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Hillier L.W., et al. *Nature* 434:724-731(2005).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.