

C10orf12 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13779**Specification**

C10orf12 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8N655
Other Accession	NM_015652 , NP_056467
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Pig, Horse, Bovine, Guinea Pig
Clonality	Rabbit
Calculated MW	Polyclonal 137kDa KDa

C10orf12 antibody - C-terminal region - Additional Information

Alias Symbol	DKFZp564P1916, FLJ13022
Other Names	
Uncharacterized protein C10orf12, C10orf12	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C10orf12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C10orf12 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


C10orf12 antibody - C-terminal region - Protein Information**C10orf12 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

C10orf12 antibody - C-terminal region - Images



WB Suggested Anti-C10orf12 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: Human kidney

C10orf12 antibody - C-terminal region - References

- Deloukas P.,et al.Nature 429:375-381(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Matsuoka S.,et al.Science 316:1160-1166(2007).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).