

**TMEM146 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13814****Specification**

---

**TMEM146 antibody - N-terminal region - Product Information**

|                   |                                                       |
|-------------------|-------------------------------------------------------|
| Application       | WB                                                    |
| Primary Accession | <a href="#">Q86XM0</a>                                |
| Other Accession   | <a href="#">NM_152784</a> , <a href="#">NP_689997</a> |
| Reactivity        | Human                                                 |
| Predicted         | Human                                                 |
| Host              | Rabbit                                                |
| Clonality         | Polyclonal                                            |
| Calculated MW     | 90kDa KDa                                             |

**TMEM146 antibody - N-terminal region - Additional Information****Gene ID** 257062**Alias Symbol** **MGC39581, TMEM146****Other Names**Cation channel sperm-associated protein subunit delta, CatSper-delta, CatSperdelta,  
Transmembrane protein 146, CATSPERD, TMEM146**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-TMEM146 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

TMEM146 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**TMEM146 antibody - N-terminal region - Protein Information****Name** CATSPERD ([HGNC:28598](#))**Synonyms** TMEM146**Function**

Auxiliary component of the CatSper complex, a complex involved in sperm cell hyperactivation. Sperm cell hyperactivation is needed for sperm motility which is essential late in the preparation of sperm for fertilization. Required for CATSPER1 stability before intraflagellar transport and/or incorporation of the CatSper complex channel into the flagellar membrane.

**Cellular Location**


Cell projection, cilium, flagellum membrane {ECO:0000250|UniProtKB:E9Q9F6}; Single-pass type I membrane protein. Note=Specifically located in the principal piece of sperm tail. {ECO:0000250|UniProtKB:E9Q9F6}

### **TMEM146 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **TMEM146 antibody - N-terminal region - Images**



WB Suggested Anti-TMEM146 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

### **TMEM146 antibody - N-terminal region - References**

Ota T., et al. Nat. Genet. 36:40-45(2004).  
Grimwood J., et al. Nature 428:529-535(2004).