

RP2 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13992**Specification**

RP2 antibody - middle region - Product Information

| | |
|-------------------|---|
| Application | IHC, WB |
| Primary Accession | O75695 |
| Other Accession | NM_006003 , NP_008846 |
| Reactivity | Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog |
| Predicted Host | Human, Mouse, Rabbit, Horse, Bovine, Dog |
| Clonality | Rabbit |
| Calculated MW | Polyclonal 40kDa KDa |

RP2 antibody - middle region - Additional Information**Gene ID** 6102

| | |
|--------------|--|
| Alias Symbol | KIAA0215, TBCCD2, XRP2, NME10, DELXp11.3 |
|--------------|--|

Other Names

Protein XRP2, RP2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RP2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RP2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

RP2 antibody - middle region - Protein Information**Name** RP2**Function**

Acts as a GTPase-activating protein (GAP) involved in trafficking between the Golgi and the ciliary membrane. Involved in localization of proteins, such as NPHP3, to the cilium membrane by inducing hydrolysis of GTP ARL3, leading to the release of UNC119 (or UNC119B). Acts as a GTPase-activating protein (GAP) for tubulin in concert with tubulin-specific chaperone C, but does not enhance tubulin heterodimerization. Acts as a guanine nucleotide dissociation inhibitor towards ADP-ribosylation factor-like proteins.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cell projection, cilium. Note=Detected predominantly at the plasma membrane of rod and cone photoreceptors. Not detected in the nucleus.


Tissue Location

Ubiquitous. Expressed in the rod and cone photoreceptors, extending from the tips of the outer segment (OS) through the inner segment (IS) and outer nuclear layer (ONL) and into the synaptic terminals of the outer plexiform layer (ONL). Also detected in the bipolar, horizontal and amacrine cells in the inner nuclear layer (INL), extending to the inner plexiform layer (IPL) and through the ganglion cell layer (GCL) and into the nerve fiber layer (NFL) (at protein level).


RP2 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RP2 antibody - middle region - Images

Immunohistochemistry with Brain, cerebellum tissue at an antibody concentration of 5 μ g/ml using anti-RP2 antibody (A113992)



90 kDa__
65 kDa__
40 kDa__
31 kDa__
22 kDa__

WB Suggested Anti-RP2 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500
Positive Control: Human Muscle

RP2 antibody - middle region - References

Schwahn U.,et al.Nat. Genet. 19:327-332(1998).
Ross M.T.,et al.Nature 434:325-337(2005).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).
Chapple J.P.,et al.Hum. Mol. Genet. 9:1919-1926(2000).
Bartolini F.,et al.J. Biol. Chem. 277:14629-14634(2002).