

SLMO2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14007**Specification**

SLMO2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A5GFX0
Other Accession	NM_016045 , NP_057129
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21kDa KDa

SLMO2 antibody - N-terminal region - Additional Information**Gene ID** 100134984**Alias Symbol** C20orf45, PRELID3B, dJ543J19.5
Other Names
Protein slowmo homolog 2, SLMO2**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SLMO2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


SLMO2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLMO2 antibody - N-terminal region - Protein Information**Name** PRELID3B**Synonyms** SLMO2**SLMO2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLMO2 antibody - N-terminal region - Images



WB Suggested Anti-SLMO2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Human Stomach