

Dctn4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14018**Specification**

Dctn4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q3TOY2
Other Accession	NM_026302 , NP_080578
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

Dctn4 antibody - C-terminal region - Additional Information

Alias Symbol	1110001K06Rik, 4930547K17Rik, C130039E17Rik, p62
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Dctn4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

Dctn4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Dctn4 antibody - C-terminal region - Protein Information**Dctn4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dctn4 antibody - C-terminal region - Images

WB Suggested Anti-Dctn4 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart