

Pex2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14036**Specification**

Pex2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8C437-3
Other Accession	NM_021483 , NP_067458
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68kDa KDa

Pex2 antibody - C-terminal region - Additional Information

Alias Symbol **1700016J08Rik, PXR2, Pex2, TRIP8b**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Pex2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Pex2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Pex2 antibody - C-terminal region - Protein Information**Pex2 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Pex2 antibody - C-terminal region - Images



WB Suggested Anti-Pex2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Liver