

TTC27 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14047**Specification**

TTC27 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6P3X3
Other Accession	NM_017735 , NP_060205
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	97kDa KDa

TTC27 antibody - N-terminal region - Additional Information**Gene ID** 55622

Alias Symbol	FLJ20272
--------------	-----------------

Other Names
Tetratricopeptide repeat protein 27, TPR repeat protein 27, TTC27**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TTC27 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TTC27 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


TTC27 antibody - N-terminal region - Protein Information**Name** TTC27**TTC27 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TTC27 antibody - N-terminal region - Images



WB Suggested Anti-TTC27 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: Human Muscle

TTC27 antibody - N-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).

Hillier L.W., et al. Nature 434:724-731(2005).

Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).