

Ppp2r3c antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14052**Specification**

Ppp2r3c antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6AXZ3
Other Accession	NM_001014196 , NP_001014218
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rabbit, Chicken, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 49kDa KDa

Ppp2r3c antibody - N-terminal region - Additional Information**Gene ID** 362739**Alias Symbol** **RGD1309207****Other Names**

Serine/threonine-protein phosphatase 2A regulatory subunit B'' subunit gamma, Ppp2r3c

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ppp2r3c antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ppp2r3c antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ppp2r3c antibody - N-terminal region - Protein Information**Name** Ppp2r3c**Function**

May regulate MCM3AP phosphorylation through phosphatase recruitment. May act as a negative regulator of ABCB1 expression and function through the dephosphorylation of ABCB1 by TFPI2/PPP2R3C complex. May play a role in the activation-induced cell death of B- cells.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9JK24}. Cytoplasm {ECO:0000250|UniProtKB:Q9JK24}.


Note=Excluded from the nucleoli Localization is cell cycle-dependent. Localizes to the cytoplasm during cytokinesis. {ECO:0000250|UniProtKB:Q9JK24}

Ppp2r3c antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ppp2r3c antibody - N-terminal region - Images



WB Suggested Anti-Ppp2r3c Antibody Titration: 1.0 µg/ml
Positive Control: Rat Brain