

CENPN antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14084**Specification**

CENPN antibody - middle region - Product Information

Application	WB, IF
Primary Accession	O96H22
Other Accession	NM_001100625 , NP_001094095
Reactivity	Human, Mouse, Rat, Rabbit, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

CENPN antibody - middle region - Additional Information**Gene ID** 55839**Alias Symbol** BM039, C16orf60, CENP-N, FLJ13607, FLJ22660**Other Names**

Centromere protein N, CENP-N, Interphase centromere complex protein 32, CENPN, C16orf60, ICEN32

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CENPN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CENPN antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

CENPN antibody - middle region - Protein Information**Name** CENPN**Synonyms** C16orf60, ICEN32**Function**

Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation. The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in

incorporation of newly synthesized CENPA into centromeres. CENPN is the first protein to bind specifically to CENPA nucleosomes and the direct binding of CENPA nucleosomes by CENPN is required for centromere assembly. Required for chromosome congression and efficiently align the chromosomes on a metaphase plate.

Cellular Location


Nucleus. Chromosome, centromere, kinetochore. Note=Localizes exclusively in the kinetochore domain of centromeres Kinetochore-bound levels decrease when cells enter mitosis and increase again when cells exit mitosis

CENPN antibody - middle region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CENPN antibody - middle region - Images




Sample Type : HeLa
Primary Antibody Dilution: 4 µg/ml
Secondary Antibody : Anti-rabbit Alexa 546
Secondary Antibody Dilution: 2 µg/ml
Gene Name : CENPN



Sample Type : Huh-7
Primary Antibody Dilution: 4 $\mu\text{g/ml}$
Secondary Antibody : Anti-rabbit Alexa 546
Secondary Antibody Dilution: 2 $\mu\text{g/ml}$
Gene Name : CENPN



Sample Type : MCF7
Primary Antibody Dilution: 4 $\mu\text{g/ml}$
Secondary Antibody : Anti-rabbit Alexa 546
Secondary Antibody Dilution: 2 $\mu\text{g/ml}$
Gene Name : CENPN



WB Suggested Anti-CENPN Antibody Titration: 0.2-1 $\mu\text{g/ml}$
Positive Control: Jurkat cell lysate
CENPN is supported by BioGPS gene expression data to be expressed in Jurkat