

PLSCR1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14134

Specification

PLSCR1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O15162
Other Accession	NM_021105 , NP_066928
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

PLSCR1 antibody - N-terminal region - Additional Information

Gene ID 5359

Alias Symbol **MMTRA1B**

Other Names

Phospholipid scramblase 1, PL scramblase 1, Ca(2+)-dependent phospholipid scramblase 1, Erythrocyte phospholipid scramblase, MmTRA1b, PLSCR1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PLSCR1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

PLSCR1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PLSCR1 antibody - N-terminal region - Protein Information

Name PLSCR1

Function

Catalyzes calcium-induced ATP-independent rapid bidirectional and non-specific movement of phospholipids (lipid scrambling or lipid flip-flop) between the inner and outer leaflet of the plasma membrane resulting in collapse of the phospholipid asymmetry which leads to phosphatidylserine externalization on the cell surface (PubMed:10770950, PubMed:18629440, PubMed:23590222, PubMed:23659204, PubMed:<a href="http://www.uniprot.org/citations/24343571"

target="_blank">>24343571, PubMed:>24648509, PubMed:>29748552, PubMed:>32110987, PubMed:>8663431, PubMed:>9218461, PubMed:>9485382, PubMed:>9572851). Mediates calcium-dependent phosphatidylserine externalization and apoptosis in neurons via its association with TRPC5 (By similarity). Also exhibits magnesium-dependent nuclease activity against double-stranded DNA and RNA but not single-stranded DNA and can enhance DNA decatenation mediated by TOP2A (PubMed:>17567603, PubMed:>27206388). Negatively regulates FcR-mediated phagocytosis in differentiated macrophages (PubMed:>26745724). May contribute to cytokine-regulated cell proliferation and differentiation (By similarity). May play a role in the antiviral response of interferon (IFN) by amplifying and enhancing the IFN response through increased expression of select subset of potent antiviral genes (PubMed:>15308695). Inhibits the functions of viral transactivators, including human T-cell leukemia virus (HTLV)-1 protein Tax, human immunodeficiency virus (HIV)-1 Tat, human hepatitis B virus (HBV) HBx, Epstein-Barr virus (EBV) BZLF1 and human cytomegalovirus IE1 and IE2 proteins through direct interactions (PubMed:>22789739, PubMed:>23501106, PubMed:>25365352, PubMed:>31434743, PubMed:>35138119). Also mediates the inhibition of influenza virus infection by preventing nuclear import of the viral nucleoprotein/NP (PubMed:>29352288, PubMed:>35595813). Plays a crucial role as a defense factor against SARS-CoV-2 independently of its scramblase activity by directly targeting nascent viral vesicles to prevent virus-membrane fusion and the release of viral RNA into the host-cell cytosol (PubMed:>37438530).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell membrane; Lipid-anchor; Cytoplasmic side. Nucleus. Cytoplasm. Cytoplasm, perinuclear region Note=Localizes to the perinuclear region in the presence of RELT (PubMed:22052202). Palmitoylation regulates its localization to the cell membrane or the nucleus; trafficking to the cell membrane is dependent upon palmitoylation whereas in the absence of palmitoylation, localizes to the nucleus (PubMed:12564925)

Tissue Location


Expressed in platelets, erythrocyte membranes, lymphocytes, spleen, thymus, prostate, testis, uterus, intestine, colon, heart, placenta, lung, liver, kidney and pancreas. Not detected in brain and skeletal muscle.

PLSCR1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)


- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLSCR1 antibody - N-terminal region - Images

WB Suggested Anti-PLSCR1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: Hela cell lysate




Host: Rabbit

Target Name: PLSCR1

Sample Tissue: Human 721_B

Antibody Dilution: 1.0µg/ml PLSCR1 is supported by BioGPS gene expression data to be expressed in 721_B



Host: Rabbit

Target Name: PLSCR1

Sample Tissue: Human 293T

Antibody Dilution: 1.0 μ g/mlPLSCR1 is strongly supported by BioGPS gene expression data to be expressed in Human HEK293T cells

PLSCR1 antibody - N-terminal region - References

- Zhou Q.,et al.J. Biol. Chem. 272:18240-18244(1997).
Kasukabe T.,et al.Biochem. Biophys. Res. Commun. 249:449-455(1998).
Wiedmer T.,et al.Biochim. Biophys. Acta 1467:244-253(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Muzny D.M.,et al.Nature 440:1194-1198(2006).