

Lhpp Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14155**Specification**

Lhpp Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q9D7I5
Other Accession	NM_029609 , NP_083885
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30kDa KDa

Lhpp Antibody - C-terminal region - Additional Information**Gene ID** 76429**Alias Symbol** 2310007H09Rik**Other Names**

Phospholysine phosphohistidine inorganic pyrophosphate phosphatase, 3.1.3.-, 3.6.1.1, Lhpp

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Lhpp antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Lhpp Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Lhpp Antibody - C-terminal region - Protein Information**Name** Lhpp**Function**

Phosphatase that hydrolyzes imidodiphosphate, 3- phosphohistidine and 6-phospholysine. Has broad substrate specificity and can also hydrolyze inorganic diphosphate, but with lower efficiency (By similarity).

Cellular Location


Cytoplasm. Nucleus.

Lhpp Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lhpp Antibody - C-terminal region - Images



WB Suggested Anti-Lhpp Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart

Lhpp Antibody - C-terminal region - References

Carninci P., et al. Science 309:1559-1563(2005).