

PARVG Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14157**Specification**

PARVG Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q9HBI0
Other Accession	XP_005261759
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36kDa kDa

PARVG Antibody - C-terminal region - Additional Information**Gene ID** 64098

Alias Symbol	PARVG,
Other Names	
Gamma-parvin, PARVG	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-PARVG antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

PARVG Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PARVG Antibody - C-terminal region - Protein Information**Name** PARVG**Function**

Probably plays a role in the regulation of cell adhesion and cytoskeleton organization.

Cellular Location

Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton Note=Constituent of focal adhesions.

Tissue Location

Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow


and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in tumors, but at a lower level than in the corresponding normal tissues.

PARVG Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PARVG Antibody - C-terminal region - Images



Host: Rabbit
Target Name: PARVG
Sample Tissue: Large intestine Tumor lysates
Antibody Dilution: 1µg/ml

PARVG Antibody - C-terminal region - References

Oliski T.M.,et al.J. Cell Sci. 114:525-538(2001).
Collins J.E.,et al.Genome Res. 13:27-36(2003).
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham I.,et al.Nature 402:489-495(1999).