

### Grem1 Antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al14165

### **Specification**

# **Grem1 Antibody - middle region - Product Information**

Application WB
Primary Accession 035793

Other Accession NM 019282, NP 062155

Reactivity Human, Mouse, Rat, Rabbit, Goat, Horse,

Bovine, Guinea Pig, Dog

Predicted Human, Mouse, Rat, Rabbit, Pig, Goat,

Horse, Bovine, Guinea Pig, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 20kDa KDa

# Grem1 Antibody - middle region - Additional Information

**Gene ID 50566** 

Alias Symbol Cktsf1b1, drm

**Other Names** 

Gremlin-1, Cysteine knot superfamily 1, BMP antagonist 1, Down-regulated in Mos-transformed cells protein, Grem1, Cktsf1b1, Drm

#### **Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

#### **Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Grem1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

### **Precautions**

Grem1 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

#### Grem1 Antibody - middle region - Protein Information

#### Name Grem1

Synonyms Cktsf1b1, Drm

#### **Function**

Cytokine that may play an important role during carcinogenesis and metanephric kidney organogenesis, as a BMP antagonist required for early limb outgrowth and patterning in maintaining the FGF4-SHH feedback loop. Down-regulates the BMP4 signaling in a dose-dependent manner (By similarity). Antagonist of BMP2; inhibits BMP2-mediated differentiation of osteoblasts



(in vitro) (By similarity). Acts as inhibitor of monocyte chemotaxis (PubMed:<a href="http://www.uniprot.org/citations/15528323" target="\_blank">15528323</a>). Can inhibit the growth or viability of normal cells but not transformed cells when is overexpressed.

## **Cellular Location**

Secreted.

#### **Tissue Location**


Highly expressed in the brain, kidney, spleen, and testis and weakly expressed in the lung and liver. Predominantly expressed in differentiated cells as neurons in brain, type I cells in lung and globlet cells in intestine.

# Grem1 Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Grem1 Antibody - middle region - Images



Host: Rabbit

Target Name: Grem1

Sample Tissue: Rat Testis lysates Antibody Dilution: 1.0µg/ml

# Grem1 Antibody - middle region - References

Topol L.Z., et al. Mol. Cell. Biol. 17:4801-4810(1997). Chen B., et al. J. Immunol. 173:5914-5917(2004).