

Tekt3 Antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14176**Specification**

Tekt3 Antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q6X6Z7
Other Accession	NM_027660 , NP_081936
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

Tekt3 Antibody - N-terminal region - Additional Information**Gene ID** 71062

Alias Symbol	4933407G07Rik
Other Names	
Tektin-3, Tekt3	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Tekt3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Tekt3 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Tekt3 Antibody - N-terminal region - Protein Information**Name** Tekt3**Function**

Microtubule inner protein (MIP) part of the dynein-decorated doublet microtubules (DMTs) in cilia and flagellar axoneme. Forms filamentous polymers in the walls of ciliary and flagellar microtubules (By similarity). Required for normal sperm mobility (PubMed:18951373).

Cellular Location

Cytoplasm, cytoskeleton, cilium axoneme {ECO:0000250|UniProtKB:A6H782}. Cell projection,

cilium, flagellum {ECO:0000250|UniProtKB:A6H782, ECO:0000250|UniProtKB:Q4V8G8}
Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane
{ECO:0000250|UniProtKB:A6H782, ECO:0000250|UniProtKB:Q4V8G8}; Peripheral membrane
protein {ECO:0000250|UniProtKB:Q4V8G8}. Note=In spermatozoa, preferentially localizes to the
flagella, but also found in the head (By similarity). In the sperm flagellum, localizes to the
periaxonemal region where it associates with the mitochondrial sheath and outer dense fibers (By
similarity). Not detected in the central axonemal region of the flagellum (By similarity). Associates
with the acrosome membrane in the equatorial segment of the sperm head (By similarity). Also
detected just below the plasma membrane in the post- acrosomal region where it might localize to
the postacrosomal dense lamina (By similarity). However, other studies report little or no
expression in the postacrosomal region (By similarity). Translocates from the postacrosomal
region to the equatorial segment after sperm activation (By similarity). Retained in the
postacrosomal region, but not the equatorial segment, following the acrosome reaction (By
similarity). Some studies report strong expression in the anterior cap region (By similarity).
However, other studies report little or no expression in the acrosomal cap (By similarity)
{ECO:0000250|UniProtKB:A6H782, ECO:0000250|UniProtKB:Q4V8G8,
ECO:0000250|UniProtKB:Q9BXF9}

Tissue Location


Expressed preferentially in testis. Expressed predominantly in late pachytene spermatocytes and
early round spermatids (PubMed:14735490). Expressed in spermatozoa (PubMed:36708031).

Tekt3 Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tekt3 Antibody - N-terminal region - Images



Host: Rabbit
Target Name: Tekt3
Sample Tissue: Mouse Muscle lysates
Antibody Dilution: 1.0µg/ml

Tekt3 Antibody - N-terminal region - References

Roy A.,et al.Mol. Reprod. Dev. 67:295-302(2004).
Roy A.,et al.Mol. Reprod. Dev. 76:453-459(2009).