

Ppp4r4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14183**Specification**

Ppp4r4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8C0Y0
Other Accession	NM_028980 , NP_083256
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Mouse, Rabbit, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96kDa KDa

Ppp4r4 antibody - C-terminal region - Additional Information**Gene ID** 74521**Alias Symbol** 8430415E04Rik, mKIAA1622**Other Names**

Serine/threonine-protein phosphatase 4 regulatory subunit 4, Ppp4r4, Kiaa1622, Pp4r4

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Ppp4r4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Ppp4r4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ppp4r4 antibody - C-terminal region - Protein Information**Name** Ppp4r4**Synonyms** Kiaa1622, Pp4r4**Function**

Putative regulatory subunit of serine/threonine-protein phosphatase 4.

Cellular Location


Cytoplasm.

Ppp4r4 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ppp4r4 antibody - C-terminal region - Images



WB Suggested Anti-Ppp4r4 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Liver

Ppp4r4 antibody - C-terminal region - References

Okazaki N., et al. DNA Res. 11:205-218(2004).
Carninci P., et al. Science 309:1559-1563(2005).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).