

Gtf2b antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14241**Specification**

Gtf2b antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P62916
Other Accession	NM_031041 , NP_112303
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rabbit, Pig, Chicken, Horse, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35kDa KDa

Gtf2b antibody - N-terminal region - Additional Information**Gene ID** 81673**Other Names**

Transcription initiation factor IIB, General transcription factor TFIIB, RNA polymerase II alpha initiation factor, Gtf2b

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Gtf2b antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Gtf2b antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Gtf2b antibody - N-terminal region - Protein Information**Name** Gtf2b {ECO:0000312|RGD:619735}**Function**

General transcription factor that plays a role in transcription initiation by RNA polymerase II (Pol II). Involved in the pre-initiation complex (PIC) formation and Pol II recruitment at promoter DNA. Together with the TATA box-bound TBP forms the core initiation complex and provides a bridge between TBP and the Pol II- TFIIF complex. Released from the PIC early following the onset of transcription during the initiation and elongation transition and reassociates with TBP during the next transcription cycle. Associates with chromatin to core promoter-specific regions. Binds to two distinct DNA core promoter consensus sequence elements in a TBP-independent manner; these

IIB-recognition elements (BREs) are localized immediately upstream (BREu), 5'-[GC][GC][GA]CGCC-3', and downstream (BREd), 5'- [GA]T[TGA][TG][GT][TG][TG]-3', of the TATA box element. Modulates transcription start site selection. Exhibits also autoacetyltransferase activity that contributes to the activated transcription.

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q00403}. Chromosome {ECO:0000250|UniProtKB:Q00403}. Note=Non-acetylated form colocalizes with DNA in the G0/1, S and G2 phases of the cell cycle, but not during mitosis. Acetylated form colocalizes at transcriptionally silent mitotic chromatids during mitosis at metaphase, anaphase, and telophase phases of the cell cycle {ECO:0000250|UniProtKB:Q00403}

Gtf2b antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Gtf2b antibody - N-terminal region - Images



WB Suggested Anti-Gtf2b Antibody Titration: 1.0 µg/ml
Positive Control: Rat lung

Gtf2b antibody - N-terminal region - References

Tsuboi A., et al. Nucleic Acids Res. 20:3250-3250(1992).