

**Skil antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14252****Specification**

---

**Skil antibody - C-terminal region - Product Information**

|                   |                                                                         |
|-------------------|-------------------------------------------------------------------------|
| Application       | WB                                                                      |
| Primary Accession | <a href="#">Q3TB81</a>                                                  |
| Other Accession   | <a href="#">NM_001039090</a> , <a href="#">NP_001034179</a>             |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog          |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                                  |
| Clonality         | Polyclonal                                                              |
| Calculated MW     | 71kDa kDa                                                               |

**Skil antibody - C-terminal region - Additional Information**

|              |                                                |
|--------------|------------------------------------------------|
| Alias Symbol | 9130011J04Rik, Skir, SnoN, SnoN2, sno, sno-dE3 |
|--------------|------------------------------------------------|

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Skil antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


**Precautions**

Skil antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Skil antibody - C-terminal region - Protein Information****Skil antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Skil antibody - C-terminal region - Images**

WB Suggested Anti-Skil Antibody Titration: 1.0 µg/ml  
Positive Control: Mouse Spleen