

DLD antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14295**Specification**

DLD antibody - middle region - Product Information

Application	WB, IHC
Primary Accession	B2R5X0
Other Accession	NM_000108 , NP_000099
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56kDa KDa

DLD antibody - middle region - Additional Information

Alias Symbol **DLDH, E3, GCSL, LAD, PHE3**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DLD antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DLD antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

DLD antibody - middle region - Protein Information**DLD antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DLD antibody - middle region - Images



Rabbit Anti-DLD Antibody

Catalog Number: AI14295

Formalin Fixed Paraffin Embedded Tissue: Human Pineal Tissue Observed Staining: Cytoplasmic in cell bodies and processes of pinealocytes

Primary Antibody

Concentration: 1:100


Secondary Antibody: Donkey anti-Rabbit-Cy3

Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 - 2.0 sec



WB Suggested Anti-DLD Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

DLD is supported by BioGPS gene expression data to be expressed in Jurkat