

NCF4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14338**Specification**

NCF4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A8K4F9
Other Accession	NM_013416 , NP_038202
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39kDa KDa

NCF4 antibody - C-terminal region - Additional Information

Alias Symbol **MGC3810, NCF, P40PHOX, SH3PXD4**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NCF4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

NCF4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NCF4 antibody - C-terminal region - Protein Information**NCF4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCF4 antibody - C-terminal region - Images



90 kDa__
65 kDa__
40 kDa__
31 kDa__
22 kDa__

WB Suggested Anti-NCF4 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:1562500

Positive Control: Human Muscle