

UBL4A antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al14388

# Specification

# UBL4A antibody - middle region - Product Information

Application Primary Accession Other Accession Reactivity

Predicted

Host Clonality Calculated MW WB <u>B7NZO9</u> <u>NM\_014235</u>, <u>NP\_055050</u> Human, Mouse, Rat, Rabbit, Horse, Yeast, Bovine, Guinea Pig, Dog Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog Rabbit Polyclonal 18kDa KDa

## UBL4A antibody - middle region - Additional Information

Gene ID 100328764

Alias Symbol

DX254E, DXS254E, G6PD, GDX, UBL4, GET5, MDY2, TMA24

**Other Names** Ubiquitin-like protein 4A, UBL4A

### Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

### **Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-UBL4A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions** UBL4A antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

### UBL4A antibody - middle region - Protein Information

Name UBL4A

### Function

As part of a cytosolic protein quality control complex, the BAG6/BAT3 complex, maintains misfolded and hydrophobic patches- containing proteins in a soluble state and participates in their proper delivery to the endoplasmic reticulum or alternatively can promote their sorting to the proteasome where they undergo degradation. The BAG6/BAT3 complex is involved in the post-translational delivery of tail-anchored/type II transmembrane proteins to the endoplasmic reticulum membrane. Recruited to ribosomes, it interacts with the transmembrane region of newly



synthesized tail-anchored proteins and together with SGTA and ASNA1 mediates their delivery to the endoplasmic reticulum. Client proteins that cannot be properly delivered to the endoplasmic reticulum are ubiquitinated and sorted to the proteasome. Similarly, the BAG6/BAT3 complex also functions as a sorting platform for proteins of the secretory pathway that are mislocalized to the cytosol either delivering them to the proteasome for degradation or to the endoplasmic reticulum. The BAG6/BAT3 complex also plays a role in the endoplasmic reticulum-associated degradation (ERAD), a quality control mechanism that eliminates unwanted proteins of the endoplasmic reticulum through their retrotranslocation to the cytosol and their targeting to the proteasome. It maintains these retrotranslocated proteins in an unfolded yet soluble state condition in the cytosol to ensure their proper delivery to the proteasome.

Cellular Location Cytoplasm, cytosol {ECO:0000250|UniProtKB:P11441}. Nucleus {ECO:0000250|UniProtKB:P11441}

### **UBL4A** antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### **UBL4A** antibody - middle region - Images



WB Suggested Anti-UBL4A Antibody Titration: 0.2-1 µg/ml ELISA Titer: 1:312500 Positive Control: Jurkat cell lysate

### UBL4A antibody - middle region - References

Antonellis A., et al. Submitted (DEC-2008) to the EMBL/GenBank/DDBJ databases.