

Hcrt Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14411**Specification****Hcrt Antibody - middle region - Product Information**

Application	WB
Primary Accession	O55232
Other Accession	NM_013179 , NP_037311
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14kDa KDa

Hcrt Antibody - middle region - Additional Information**Gene ID 25723**

Alias Symbol	orexin-A
Other Names	
Orexin, Hypocretin, Hcrt, Orexin-A, Hypocretin-1, Hcrt1, Orexin-B, Hypocretin-2, Hcrt2, Hcrt, Ox, Ppox	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Hcrt antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Hcrt Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Hcrt Antibody - middle region - Protein Information

Name Hcrt {ECO:0000312|RGD:2786}

Synonyms Ox, Ppox

Function

Neuropeptides that play a significant role in the regulation of food intake and sleep-wakefulness, possibly by coordinating the complex behavioral and physiologic responses of these complementary homeostatic functions (PubMed:11283317, PubMed:>11340621, PubMed:>9491897). A broader role in the homeostatic regulation of energy metabolism, autonomic function, hormonal balance and the regulation of body fluids, is also suggested (PubMed:>9491897). A modulation effect on luteinizing hormone-releasing hormone (LHRH) secretion also suggests a more minor contribution to the regulation of reproductive function (PubMed:>11340621, PubMed:>9879756).

Cellular Location

Rough endoplasmic reticulum. Cytoplasmic vesicle. Synapse Note=Associated with perikaryal rough endoplasmic reticulum as well as cytoplasmic large granular vesicles at synapses

Tissue Location


Mainly expressed in the brain, with expression in neurons in the dorsal lateral hypothalamic area (at protein level) (PubMed:9419374). Produced by a small group of neurons restricted to the lateral and posterior hypothalamus and perifornical areas (PubMed:9491897). Positive neurons project widely throughout the entire neuroaxis (PubMed:9491897). Particularly abundant projections in the cerebral cortex, olfactory bulb, hippocampus, amygdala, septum, diagonal band of Broca, bed nucleus of the stria terminalis, thalamus, anterior and posterior hypothalamus, midbrain, brainstem, and spinal cord (PubMed:9491897). Also expressed in the enteric nervous system and pancreas (PubMed:9491897). In small amount, also detected in the testis (PubMed:9491897).

Hcrt Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hcrt Antibody - middle region - Images



Host: Rabbit

Target Name: Hcrt

Sample Tissue: Rat Testis lysates

Antibody Dilution: 1.0µg/ml

Hcrt Antibody - middle region - References

- Sakurai T.,et al.Cell 92:573-585(1998).
de Lecea L.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:322-327(1998).
Hungs M.,et al.Bioessays 23:397-408(2001).
Willie J.T.,et al.Annu. Rev. Neurosci. 24:429-458(2001).