

GRP antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14417**Specification**

GRP antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P07492-3
Other Accession	NM_001012512 , NP_001012530
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15kDa KDa

GRP antibody - C-terminal region - Additional Information

Alias Symbol **BN, GRP-10, preproGRP, proGRP**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-GRP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


GRP antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

GRP antibody - C-terminal region - Protein Information**GRP antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRP antibody - C-terminal region - Images



WB Suggested Anti-GRP Antibody Titration: 1.0 μ g/ml
Positive Control: HepG2 Whole Cell