

OR1L8 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14442**Specification**

OR1L8 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NGR8
Other Accession	NM_001004454 , NP_001004454
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human
Clonality	Rabbit
Calculated MW	Polyclonal 35kDa kDa

OR1L8 antibody - C-terminal region - Additional Information**Gene ID** 138881**Alias Symbol** **OR9-24**
Other Names
Olfactory receptor 1L8, Olfactory receptor OR9-24, OR1L8**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-OR1L8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

OR1L8 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

OR1L8 antibody - C-terminal region - Protein Information**Name** OR1L8**Function**

Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR1L8 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR1L8 antibody - C-terminal region - Images



WB Suggested Anti-OR1L8 Antibody Titration: 1.0 µg/ml

Positive Control: Fetal Muscle

OR1L8 antibody - C-terminal region - References

Suwa M., et al. Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.

Humphray S.J., et al. Nature 429:369-374(2004).

Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Fuchs T., et al. Genomics 80:295-302(2002).

Malnic B., et al. Proc. Natl. Acad. Sci. U.S.A. 101:2584-2589(2004).