

F8 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14494**Specification**

F8 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P00451
Other Accession	NM_000132 , NP_000123
Reactivity	Human, Rabbit, Sheep, Horse, Bovine, Dog
Predicted	Human, Rabbit, Sheep, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79kDa kDa

F8 antibody - C-terminal region - Additional Information**Gene ID 2157**

Alias Symbol **AHF, DXS1253E, F8B, F8C, FVIII, HEMA**

Other Names

Coagulation factor VIII, Antihemophilic factor, AHF, Procoagulant component, Factor VIIIa heavy chain, 200 kDa isoform, Factor VIIIa heavy chain, 92 kDa isoform, Factor VIII B chain, Factor VIIIa light chain, F8, F8C

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-F8 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

F8 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

F8 antibody - C-terminal region - Protein Information

Name F8

Synonyms F8C

Function

Factor VIII, along with calcium and phospholipid, acts as a cofactor for F9/factor IXa when it converts F10/factor X to the activated form, factor Xa.

Cellular Location


Secreted, extracellular space.

F8 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F8 antibody - C-terminal region - Images



Host: Rabbit

Target Name:F8

Sample Tissue:721_B Whole Cell

Lane A: Primary Antibody

Lane B: Primary Antibody + Blocking Peptide

Primary Antibody

Concentration:1µg/ml

Peptide Concentration: 5µg/ml

Lysate Quantity: 25ug/lane Gel

Concentration: 0.12%

F8 antibody - C-terminal region - References

Truett M.A.,et al.DNA 4:333-349(1985).

Wood W.I.,et al.Nature 312:330-337(1984).

Levinson B.,et al.Genomics 14:585-589(1992).

Toole J.J.,et al.Nature 312:342-347(1984).

Gitschier J.,et al.Hum. Mol. Genet. 1:199-200(1992).